

Characteristics:

Maturity	6,3
Colour of skin	G
Color of flesh	6,8
Shape of tuber	LLO
Depth of the eyes	7,5
Regularity shape of tuber	7,1
Tuber length (50-60 mm)	9,5 cm
Average size (mm)	53
Dry matter percentage	22,8
Suitability for cooking	8 B
Dormancy	6,2
Fusarium	1

Comments:

9 = positive 1 = negative

Yellowness of flesh: 8 = yellow 4 = white

Colour of skin: L=light, D=dark, W=white, G=yellow, R=red

Shape of tuber: R = Round O = Oval L = Long

Cooking type: A = Firm D = Very mealy

Resistance:

Common scab	7,1
Bruise index (0-50)	7
Secondary growth	7
Yntn-virus	7
Foliage blight	4
Ro1	9
Ro2/3	8
Ro5	9
Pa2	8
Pa3	2
wz 1 (D1)	10
wz 2 (G1)	
wz 6 (O1)	4
wz 18 (T1)	8

Bruise index (0=free - 50=very heavy)

RESISTANCE TO GOLDEN NEMATODE:

Ro1(=A), Ro2/Ro3(=BC), Ro4(=A), Ro5(=F), Pa2(=D), Pa3(=E)

(9 = high resistant; 1 = very susceptible)


WART DISEASE:

wz 1(D1), wz 2/6 (G1/O1), wz 18 (T1)

(10 = resistant, 3 = very susceptible)

Fertilisation (kg/ha):

depending on the soil reserve and applicable national legislation.



To prevent mineral deficiencies, apply these elements in time (B, Mg, Mn, S)


Planting distance:

Size	Rowspace		Plants/hectare (x 1000)	
	75 cm	90 cm	75 cm	90 cm
28/35	33	28	40,4	39,7
35/45	40	33	33,3	33,7
45/50	42	35	31,7	31,7
45/50 CUT	35	29	38,1	38,3
50/55 CUT	38	32	35,1	34,7
50/55	44	37	30,3	30,0
55/60	44	37	30,3	30,0

Pay attention to:

- Do not plant early in cold soil (> 8° C)
- Avoid too strong germination
- Do not damage sprouts mechanical or chemical
- Do not plant deep.
- Lady Anna grows deep in the row
- After harvest, dry the potatoes by ventilation

Storage:



6 °C
8 months